arXiv:1607.01500v2 [math.NT] 3 Dec 2016

General proof for irrationality of infinite sums based on Fourier's proof

Tomer Shushi

Department of Physics, Ben-Gurion University of the Negev, Beersheba 8410501, Israel

December 6, 2016

Abstract

In this paper we review a general proof for the irrationality property of numbers which take a certain form of infinite sums.

Keywords: Fourier's irrationality proof, Infinite sum numbers, lr-rational numbers

Introduction:

Proofs for irrationality are sometimes very difficult to obtain, for instance the proofs of the irrationality of π/e or $\pi+e$ as well as $\pi-e$ are open problems in mathematics. The proofs for the irrationality of different numbers are sometimes based on different techniques and approaches (see, for instance, [1,2,3,4]). The origins of the proof in this paper seems to be unknown.

Define a number P_χ as the infinite sum

$$P_\chi = \sum_{n=0}^{\infty} \frac{\chi(n)}{n!}.$$

For any $n = 0, 1, ...,$ $\chi(n)$ is a positive integer, and $0 \leq \chi(n) \leq M$ where M is a known number (for instance $M = 100$ so $0 \leq \chi(n) \leq 100$ $\forall n = 0, 1, ...$). Note that for finite M, P_χ is a finite number, since

$$\sum_{n=0}^{\infty} \frac{\chi(n)}{n!} \leq M \sum_{n=0}^{\infty} \frac{1}{n!} = M \cdot e.$$

Theorem 1 *P_χ is irrational number if the assumption that this number is rational implies that its denominator should be greater than M.*

Proof. To prove the Theorem we first assume that P_χ is rational, i.e. P_χ can be written as the division between two integers,

$$P_\chi = \sum_{n=0}^{\infty} \frac{\chi(n)}{n!} = \frac{a}{b}.$$

Now we define a number X

$$X := b! \left(\sum_{n=0}^{\infty} \frac{\chi(n)}{n!} - \sum_{n=0}^{b} \frac{\chi(n)}{n!} \right) = b! \sum_{n=b+1}^{\infty} \frac{\chi(n)}{n!} > 0, \tag{1}$$

and recall that we assumed that P_χ is a rational number a/b, therefore we argue that X can also get only non-negative integer numbers since

$$\begin{aligned} X &= b! \left(\sum_{n=0}^{\infty} \frac{\chi(n)}{n!} - \sum_{n=0}^{b} \frac{\chi(n)}{n!} \right) = \frac{ab!}{b} - \sum_{n=0}^{b} \frac{\chi(n) b!}{n!} \\ &= a(b-1)! - \sum_{n=0}^{b} \frac{\chi(n) b!}{n!} \geq 1. \end{aligned}$$

Furthermore, the following inequality is well known

$$\frac{b!}{n!} < \frac{1}{(1+b)^{n-b}}, \ n \geq b+1,$$

and because $\chi(n) \leq M, \forall n \geq 1$ it is also true that

$$\frac{b!\chi(n)}{n!} < \frac{M}{(1+b)^{n-b}}, \ n \geq b+1. \tag{2}$$

Substituting (2) in (1) we get the inequality

$$\begin{aligned} X &= \sum_{n=b+1}^{\infty} \frac{b!\chi(n)}{n!} < M \sum_{n=b+1}^{\infty} \frac{1}{(1+b)^{n-b}} = M \sum_{n=1}^{\infty} \frac{1}{(1+b)^n} \\ &= M \left(\frac{1}{1 - 1/(1+b)} \right) \frac{1}{b+1} = \frac{M}{b}, \end{aligned}$$

therefore for $b > M$ it is clear that

$$X < 1,$$

and we get a contradiction since $X \geq 1$. Therefore P_χ is an irrational number. ■

Example 1 *Suppose that* P_χ *takes the form*

$$P_\chi = \frac{3}{0!} + \frac{5}{1!} + \frac{7}{2!} + \frac{3}{3!} + \frac{5}{4!} + ... \ ,$$

where $\chi(n)$ *is periodic with the values* $3, 5,$ *and* 7. *Since in this case* b *should be greater than* 7 *(we can simply check it!) then* P_χ *is an irrational number.*

Example 2 *Any number* $P_\chi - e$ *where* $0 < \chi(n) \leq M \ \forall n \geq 1$, $P_\chi - e > 0$ *is an irrational number if* P_χ *is an irrational number, proved by the previous Theorem.*

Example 3 *For a periodic sequence* P_k *of prime numbers from* $p_1 = 2$ *to* $p_d = 101$, *we can write the following number*

$$P_\chi = \frac{2}{0!} + \frac{3}{1!} + \frac{5}{2!} + \frac{7}{3!} + \frac{11}{4!} + ... + \frac{101}{5!} + \frac{97}{6!} + \frac{89}{7!} + ... \ .$$

This periodic sum is an irrational number since it follows the Theorem.

Example 4 *The following number*

$$\sum_{n=0}^{\infty} \frac{\cos(n\pi)}{n!} + e$$

is also irrational since $\cos(j\pi)$, $j = 1, 2, ...,$ *can get the values* -1 *or* 1 *and therefore in this case* $\chi(j) = 0, 2$ *which implies on the irrationality of this number.*

References

[1] Cohn, H. (2006). A Short Proof of the Simple Continued Fraction Expansion of e. The American Mathematical Monthly, 57-62.

[2] Niven, I. (2005). Irrational numbers (No. 11). Cambridge University Press.

[3] Shallit, J. (1979). Simple continued fractions for some irrational numbers. Journal of Number Theory, 11(2), 209-217.

[4] Zudilin, W. W. (2001). One of the numbers $\zeta(5)$, $\zeta(7)$, $\zeta(9)$, $\zeta(11)$ is irrational. Russian Mathematical Surveys, 56(4), 774-776.